

## Patient Information – Please attach a copy of the patient’s insurance card

Patient Name:		Date of Birth:	<input type="checkbox"/> Male <input type="checkbox"/> Female	
Address:		City:	State:	Zip:
Phone Number:	Alternate Phone Number:		Language:	
<b>Allergies (Required):</b>		<input type="checkbox"/> NKDA	Height:	Weight:
				SSN:
<b>Product Shipping Options:</b> <input type="checkbox"/> Patient’s Home <input type="checkbox"/> Prescriber Office <input type="checkbox"/> Alternative Address:				

## Prescriber Information

Practice Name:		Office Contact:		
Prescriber:		NPI:	DEA:	
Practice Address:		City:	State:	Zip:
Phone Number:		Fax Number:		

## Clinical Information – Please send all available chart notes including lab results

Diagnosis/ICD-10:				
BMD/T-score:		Is the patient new to therapy? <input type="checkbox"/> Yes <input type="checkbox"/> No	Does the patient have a history of osteoporotic fracture? <input type="checkbox"/> Yes <input type="checkbox"/> No	
Date of score:			If no, is the patient at high risk? <input type="checkbox"/> Yes <input type="checkbox"/> No	
			If yes, date of fracture:	
			Location of fracture:	
Prior Failed Medications: <input type="checkbox"/> Actonel® <input type="checkbox"/> Boniva® <input type="checkbox"/> Forteo® <input type="checkbox"/> Fosamax® <input type="checkbox"/> Prolia® <input type="checkbox"/> Reclast® <input type="checkbox"/> Other:				

## Prescription Information

Medication	Strength	Directions	Quantity	Refill
<input type="checkbox"/> Boniva®	3mg/3mL prefilled syringe	Inject the contents of 1 syringe (3 mg) intravenously every 3 months. To be administered by a healthcare professional	1 prefilled syringe	
<input type="checkbox"/> Forteo®	600 mcg/2.4 mL pen	Inject 1 dose (20 mcg) subcutaneously once daily. Discard device 28 days after first use. Dispensed with BD Mini™ Pen Needles (28 needles per 1 pen dispensed)	1 pen (4 weeks supply)	
<input type="checkbox"/> Prolia®	60 mg/1 mL prefilled syringe	Inject the contents of 1 syringe (60 mg) subcutaneously every 6 months	1 prefilled syringe	
<input type="checkbox"/> Tymlos™	2 mg/mL prefilled syringe	Inject 1 dose (80 mcg) subcutaneously once daily. Discard device 30 days after first use. Dispensed with BD Short™ Pen Needles (30 needles per 1 pen dispensed)	1 prefilled syringe	
<input type="checkbox"/> Other:				

## Prescriber Signature and Date (Please sign and date below)

Dispense as Written	Date	Substitution Permissible	Date
<input type="checkbox"/> Check here to authorize Curexa™ and its representatives to act as an agent to initiate and execute the insurance prior authorization process and receive forms on the prescriber’s behalf. “I certify that the above therapy is medically necessary and the above information is accurate to the best of my knowledge”			

**Important Notice:** This form and its contents may contain private and confidential information that is intended for the individual or entity to which it is addressed. This transmission may contain information that is exempt from disclosure under laws including but not limited to the Health Insurance Portability and Accountability Act (HIPAA). Unless explicitly stated, you are strictly prohibited from disseminating, copying or distributing any material contained within. Violators will be prosecuted to the fullest extent of the law. If you received this communication in error, please notify us immediately and destroy this form and its contents.